

**Gynae Fast Track Clinic**  
Tuesday to Friday  
01935 384 454  
07557 313 816

**Colposcopy team**  
Monday to Thursday  
01935 384 622

**Jasmine Ward**  
Weekends and evenings  
01935 384 385

**If you need this leaflet  
in another format, eg.  
large print or a different  
language, please ask a  
member of staff.**

Yeovil District Hospital  
NHS Foundation Trust  
Higher Kingston  
Yeovil  
Somerset  
BA21 4AT

# Hysteroscopy under general anaesthetic

Gynaecology department

01935 475 122  
[yeovilhospital.nhs.uk](http://yeovilhospital.nhs.uk)

Ref: 14-16-146  
Review: 04/18

01935 475 122  
[yeovilhospital.nhs.uk](http://yeovilhospital.nhs.uk)

Your operation will be done as a day case under general anaesthetic.

## What is hysteroscopy

Hysteroscopy is the technique of inspecting the cavity of the uterus (womb) under direct vision. This is done by using a hysteroscope (small telescope) which is inserted via the vagina and cervix (neck of the womb).

At the end of the hysteroscopy, we often take a small piece of tissue from the lining of the womb for examination.

Following the operation, you are likely to have some vaginal blood loss and you should wear pads not tampons until the bleeding has stopped.

Any discomfort should be relieved by taking mild pain relief medication such as Paracetamol.

Before you leave, you will be advised of any follow up arrangement.

You may return to work when you feel comfortable to do so, usually within 2-3 days.

## Complications

Complications after this type of operation are unusual. However, if you have any of the following:

- Prolonged heavy bleeding / or pass clots
- An offensive discharge
- Fever
- Severe pain

### Do get in touch with your own doctor

## You will have been advised of the following

- You need to be accompanied home by an adult after your operation
- There will be somebody with you for 24 hours after your operation
- You live within one hour's drive of the hospital
- You should not cook or operate kitchen machinery for 24 hours after your operation
- You should not drive a car, ride a bicycle, operate heavy machinery or make important legal or business decisions for 48 hours after your operation